Rylaarsdam 1983), although this is not always the case (Fitzhardinge 1985, Harriott 1985). Relative recruitment rates have been shown to vary from year to year, and among sites on the Great Barrier Reef (Wallace 1985 a).

There appears to be an inverse correlation between success of larval recruitment and propagation by fragmentation in some coral species (Kojis & Quinn 1981 b, Highsmith 1982, Wallace 1985 b). Sammarco (1985, 1987) reported that, while planular recruitment of Acropora spp. on the Great Barrier Reef is high, juvenile recruits are rare in the Caribbean and populations there are structured primarily by asexual processes. Temporal and regional variations in biotic and environmental factors can cause differences in reproductive and recruitment patterns of coral communities.

The pan-Pacific coral Pocillopora damicornis provides an example of how life history characteristics may vary with respect to local conditions (Richmond 1985, 1987 a). Enewetak Atoll and Hawaii are characterized as having low rates of predation on P. damicornis, relatively low seasonal variability, yet relatively high frequencies of mortality-causing disturbances (typhoons and winter storms). P. damicornis colonies planulate monthly throughout the year in Hawaii and the Central Pacific, and the oldest colonies observed are estimated to be less than 10 yr old (branch lengths < 20 cm). The eastern Pacific of Panama is characterized as having high rates of predation on P. damicornis, high levels of seasonal variability, and low frequencies of mortality-causing events. In this area, P. damicornis has never been found to release planulae, has a higher colony growth rate, and is the dominant coral in terms of competitive interactions with other corals, with some colonies estimated as over 70 yr old (branch lengths > 240 cm).

Theories on the evolution of life history characteristics have been proposed which suggest that under conditions of environmental instability, where lethal disturbances occur at a relatively high rate, formation of large numbers of motile propagules should occur, while under stable conditions, selection will favor clonal (asexual) growth (Williams 1975, Maynard-Smith 1978). Likewise, under conditions of low juvenile versus high adult mortality, and relatively low competitive ability versus competitive dominance, selection will favor the sexual mode (Abrahamson 1980, Douglas 1981). *Pocillopora damicornis* fits these predicted patterns over its distributional range.

RESOURCE MANAGEMENT IMPLICATIONS

As coral reefs throughout the world are showing signs of degradation, management of reef resources is becoming a growing concern. In the case of corals which spawn during a very brief period each year, the presence of contaminants such as petroleum products, pesticides, herbicides and heavy metals from soils may prevent successful fertilization of eggs by sperm, and hence, severely limit coral recruitment (Richmond unpubl.). It is suggested that pollution levels which may not affect adult coral colonies could still be responsible for the eventual loss of reefs if reproductive processes are disturbed. In Guam and in Okinawa (southern Japan), the peak coral spawning occurs during the rainy season, when levels of coastal contamination via runoff are at their highest (pers. obs.).

With reef degradation and destruction occurring on a global scale, an application of the reproductive data is in the area of reef recovery. Areas of reef which have been destroyed may be re-seeded, and the most efficient means will depend on local conditions. In areas where sedimentation is high, corallivores are present in large numbers, and/or disturbance rate is low, cementing larger numbers of smaller fragments may be more effective based on reproduction and recruitment data. In areas where environmental conditions support sexual reproduction of corals, juvenile mortality is expected to be relatively low, and suitable substratum is available, transplantation of gravid adult colonies into an area may result in highest return of effort. Reef re-seeding is an expensive process (Harriott & Fisk 1989) and management efforts might best be expended in coral reef protection and conservation.

SUMMARY

Data on the reproductive biology of 200+ scleractinian corals indicate several apparent trends. Species which broadcast spawn outnumber those which broad planula larvae. Broadcast spawners typically have limited, annual spawning periods, while most brooders are iteroparous, releasing larvae over a large part of the year. Most corals reported to date are hermaphroditic, some of which have been found to self-fertilize readily in the laboratory; the capacity for selfing is low to non-existent in other hermaphroditic species. There are reports of coral populations which are apparently sterile, particularly those at the extreme limits of their distributions.

There appear to be several geographical trends in coral reproduction. The majority of species on the Great Barrier Reef participate in an annual synchronous spawning event following full moon in the austral spring. In other areas of the Pacific, the Caribbean, and the Red Sea, there is a greater partitioning of spawning periods over more months, days, and lunar phases. Degree of synchrony among species may be related to seawater temperature ranges in each region.

Mode of reproduction within a given taxa is generally conservative, while timing may be variable within and among species. Differences in reproductive patterns may represent adaptations to local environmental conditions or may represent groups which have not been sufficiently taxonomically differentiated. A point which clearly emerges from the data reviewed in this paper is the need for examination of reproductive characteristics in conjunction with skeletal characters to elucidate what may be real problems in coral taxonomy. Do observed reproductive differences indicate that speciation has occurred among populations? If we accept the biological species definition, it is evident that morphological characters alone may fail to detect valid biological differences. As more studies of reproduction are pursued, resulting information will enable development of a more accurate understanding of the ecology and evolutionary biology of the Scleractinia.

Acknowledgements. We gratefully acknowledge UNESCO for sponsoring the meeting on Interoceanic Comparisons, held at the University of the South Pacific, Suva, Fiji in March, 1986, for which this review was originally prepared. We thank Drs C. E. Birkeland, P. L. Jokiel, R. A. Kinzie, D. A. Krupp and P. W. Sammarco for comments on manuscript drafts, R. Randall for taxonomic identifications, D. Hopper, F. Te, A. Kerr, and A. Goodis for assistance with field and laboratory examination of Guam corals, and Dr K. Yamazato for access to the corals of Okinawa. This is contribution no. 277 of the University of Guam Marine Laboratory.

LITERATURE CITED

Abe, N. (1937). Postlarval development of the coral Fungia actiniformis var. palawensis Doderlein. Palao trop. biol. Stn. Stud. 1: 73-93

Abrahamson, W. G. (1980). Demography and vegetative reproduction. In: Solbrig, O. T. (ed.) Demography and evolution in plant populations. Botanical Monographs 15. Blackwell Scientific Publication, London, p. 89–106

Andrews, J. C., Gay, S. L., Sammarco, P. W. (1989). Models of dispersal and recruitment of coral larvae around an isolated reef (Helix Reef) in the central Great Barrier Reef. Proc. 6th Int. Coral Reef Congr., Townsville (in press)

Atoda, K. (1947 a). The larva and postlarval development of some reef-building corals. I. Pocillopora damicornis (Dana). Sci. Rep. Tohoku Univ. 18: 24–47

Atoda, K. (1947b). The larva and postlarval development of some reef-building corals. II. *Stylophora pistillata* (Esper.). Sci. Rep. Tohoku Univ. 18: 48–64

Atoda, K. (1951a). The larvae and postlarval development of some reef-building corals. III. Acropora brueggemanni (Brook). J. Morph. 89: 1–13

Atoda, K. (1951b). The larva and postlarval development of some reef-building corals. IV. Galaxea aspera Quelch. J. Morph. 89: 17–35

Atoda, K. (1951c). The larva and postlarval development of some reefbuilding corals. IV. *Seriatopora hystrix* Dana. Sci. Rep. Tohoku Univ. 19: 33–39

Ayre, D. J., Resing, J. M. (1986). Sexual and asexual production of planulae in reef corals. Mar. Biol. 90: 187–190

Babcock, R. C. (1984). Reproduction and distribution of two species of *Goniastrea* (Scleractinia) from the Great Barrier Reef province. Coral Reefs 2: 187–204

Babcock, R. C. (1989). Fine-scale spatial and temporal patterns in coral recruitment. Proc. 6th Int. Coral Reef Symp.,

Townsville, in press

Babcock, R. C., Bull, G. D., Harrison, P. L., Heyward, A. J., Oliver, J. K., Wallace, C. C., Willis, B. L. (1986). Synchronous spawning of 105 scleractinian coral species on the Great Barrier Reef. Mar. Biol. 90: 379–394

Babcock, R. C., Heyward, A. J. (1986). Larval development of certain gamete-spawning scleractinian corals. Coral Reefs

5: 111-116

Baggett, L. S., Bright, T. J. (1985). Coral recruitment at the East Flower Garden Reef. Proc. 5th Int. Coral Reef Cong., Tahiti 4: 379–384

Bak, R. P. M., Engel, M. S. (1979). Distribution, abundance and survival of juvenile hermatypic corals (Scleractinia) and the importance of life history strategies in the parent coral community. Mar. Biol. 54: 341–352

Birkeland, C., Rowley, D., Randall, R. H. (1981). Coral recruitment patterns at Guam. Proc. 4th Int. Coral Reef Symp.,

Manila, 2: 339-344

Boschma, H. (1929). On the post-larval development of the coral Maendra areolata. Pap. Tortugas Lab. 26: 129–147

Bothwell, A. M. (1981). Fragmentation, a means of asexual reproduction and dispersal in the coral genus *Acropora* (Scleractinia: Astrocoeniida: Acroporidae) – a preliminary report. Proc. 4th Int. Coral Reef Symp., Manila 2: 137–144

Chornesky, E. A., Peters, E. C. (1987). Sexual reproduction and colony growth in the scleractinian coral *Porites* astreoides. Biol. Bull. mar. biol. Lab., Woods Hole 161–177

Dana, T. F. (1975). Development of contemporary eastern Pacific coral reefs. Mar. Biol. 33: 355–374

Douglas, D. A. (1981). The balance between vegetative and sexual reproduction of *Mimulus primuloides* (Scrophularíaceae) at different altitudes in California. J. Ecol. 69: 295–310

Duerden, J. E. (1902). West Indian madreporarian polyps. Mem. natn. Acad. Sci. 8: 401–597

Edmondson, C. H. (1929). Growth of Hawaiian corals. Bull. Bernice P. Bishop Mus. 58: 1–38

Edmondson, C. H. (1946). Behavior of coral planulae under altered saline and thermal conditions. Occ. Pap. Bernice P. Bishop Mus. 18: 283–304

Emery, K. O. (1962). Marine geology of Guam. Prof. Pap. U.S. Geol. Surv. 403-B: 1–76

Fadlallah, Y. H. (1983). Sexual reproduction, development and larval biology in scleractinian corals. A review. Coral Reefs 2: 129–150

Fadlallah, Y. H. (1985). Reproduction in the coral *Pocillopora verrucosa* on the reefs adjacent to the industrial city of Yanbu, Red Sea, Saudi Arabia. Proc. 5th Int. Coral Reef Symp., Tahiti 4: 314–318

Fitzhardinge, R. (1985). Spatial and temporal variability in coral recruitment in Kaneohe Bay. Proc. 5th Internat. Coral

Reef Cong., Tahiti 4: 373-378

Gilmore, M. D., Hall, B. R. (1976). Life history, growth habits, and constructional roles of Acropora cervicornis in the patch reef environment. J. sedim. Petrol. 46: 519–522

Goreau, T. H., Goreau, N. I. (1959). The physiology of skeleton formation in corals. II. Calcium deposition by hermatypic corals under various conditions in the reef. Biol. Bull. mar. biol. Lab., Woods Hole 117: 239–250

Goreau, T. H., Wells, J. M. (1967). The shallow-water scleractinia of Jamaica: revised list of species and their vertical distribution range. Bull. mar. Sci. 17: 442–452

Grigg, R. W., Wells, J. M., Wallace, C. (1981). Acropora in Hawaii, Part 1: History of the scientific record, systematics and ecology. Pacif. Sci. 35: 1-13

Hada, Y. (1932). A note of earlier stage of colony formation with the coral *Pocillopora caespitosa*. Sci. Rep. Tohuku Imp. Univ. 6: 425–431

Harrigan, J. F. (1972). The planula larva of *Pocillopora damicornis*, lunar periodicity of swarming and substratum selection behavior. Ph. D. dissertation, Univ. of Hawaii

Harriott, V. J. (1983 a). Reproductive ecology of four scleractinian species at Lizard Island, Great Barrier Reef. Coral Reefs 2: 9–18

Harriott, V. J. (1983b). Reproductive seasonality, settlement, and post-settlement mortality of *Pocillopora damicornis* (Linnaeus), at Lizard Island, Great Barrier Reef. Coral Reefs 2: 151–157

Harriott, V. J. (1985). Recruitment patterns of scleractinian corals at Lizard Island, Great Barrier Reef. Proc. 5th Int. Coral Reef Cong., Tahiti 4: 367–372

Harriott, V. J., Fisk, D. A. (1989). Coral transplantation as a reef management option. Proc. 6th Int. Coral Reef Symp., Townsville (in press)

Harrison, P. L. (1985). Sexual characteristics of scleractinian corals: systematic and evolutionary implications. Proc. 5th Int. Coral Reef Symp., Tahiti 4: 337–342

Harrison, P. L. (1989). Pseudo-gynodieocy: an unusual breeding system in the scleratinian coral Galaxea fascicularis. Proc. 6th Int. Coral Reef Symp., Townsville (in press)

Harrison, P. L., Babcock, R. C., Bull, G. D., Oliver, J. K., Wallace, C. C., Willis, B. L. (1984). Mass spawning in tropical reef corals. Science 223: 1186–1189

Heyward, A. J. (1985). Sexual reproduction in five species of the coral *Montipora*. Tech. Rep. Hawaii Inst. Mar. Biol. 37: 170–178

Heyward, A. J. (1989). Reproductive status of some Guam corals. Micronesica 21: 272–274

Heyward, A. J., Babcock, R. C. (1986). Self- and cross-fertilization in scleractinian corals. Mar. Biol. 90: 191–195

Heyward, A. J., Collins, J. D. (1985). Growth and sexual reproduction in the scleractinian coral Montipora digitata (Dana). Aust. J. mar. Freshwat. Res. 36: 441–446

Heyward, A. J., Yamazato, K., Yemin, T., Minei, M. (1987).
Sexual reproduction of corals in Okinawa. Galaxea 6: 331–343

Highsmith, R. C. (1982). Reproduction by fragmentation in corals. Mar. Ecol. Prog. Ser. 7: 207–226

Highsmith, R. C., Riggs, A. C., D'Antonio, C. M. (1980). Survival of hurricane-generated coral fragments and a disturbance model of reef calcification/growth rates. Oecologia (Berl.) 46: 322–329

Hunter, C. L. (1988). Genotypic diversity and population structure of the Hawaiian reef coral, *Porites compressa*. Ph. D. dissertation, Univ. of Hawaii

Hunter, C. L. (1989). Environmental cues controlling spawning in two Hawaiian corals: *Montipora verrucosa* and *M. dilatata*. Proc. 6th Int. Coral Reef Symp., Townsville (in press)

Jokiel, P. L. (1985). Lunar periodicity of planula release in the reef coral *Pocillopora damicornis* in relation to various environmental factors. Proc. 5th Int. Coral Reef Symp., Tahiti 4: 307-312

Jokiel, P. L. (1987). Ecology, biogeography and evolution of corals in Hawaii. Trends Ecol. Evol. 2 (7): 179–182

Jokiel, P. L., Ito, R. Y., Liu, P. M. (1985). Night irradiance and synchronization of lunar spawning of larvae in the reef coral *Pocillopora damicornis* (Linnaeus). Mar. Biol. 88: 167–174

Kawaguti, S. (1941). On the physiology of reef corals. V.

Tropisms of coral planulae considered as a factor of distribution on the reef. Palao trop. biol. Stn Stud. 2: 319–328

Kojis, B. L., Quinn, N. J. (1981 a). Aspects of sexual reproduction and larval development in the shallow water hermatypic coral Goniastrea australensis (Edwards and Haime 1857). Bull. mar. Sci. 31: 558-573

Kojis, B. L., Quinn, N. J. (1981b). Reproductive strategies in four species of *Porites* (Scleractinia). Proc. 4th Int. Coral Reef Symp., Manila 2: 145–151

Kojis, B. L., Quinn, N. J. (1982). Reproductive ecology of two Faviid corals (Coelenterata: Scleractinia). Mar. Ecol. Prog. Ser. 8: 251–255

Krupp, D. A. (1983). Sexual reproduction and early development of the solitary coral Fungia scutaria (Anthozoa: Scleractinia). Coral Reefs 2: 159–164

Loya, Y. (1976). The Red Sea coral Stylophora pistillata is an rstrategist. Nature, Lond. 259: 478–480

Marshall, S. M., Stephenson, T. A. (1933). The breeding of reef animals. Part I. The corals. Scient. Rep. Gt Barrier Reef Exped. 3: 219–245

Mavor, J. W. (1915). On the development of the coral *Agaricia* fragilis Dana. Proc. Am. Acad. Arts Sci. 51: 485–511

Maynard-Smith, J. (1978). The evolution of sex. Cambridge University Press, London

Mileikovsky, S. A. (1971). Types of larval development in marine bottom invertebrates, their distribution and ecological significance: a re-evaluation. Mar. Biol. 10: 193–213

Nakamura, S. (1984). Record of air temperature, surface water temperature and chlorinity at Sesoko in 1983. Galaxea 3:

Oliver, J., Babcock, R., Harrison, P., Willis, B. (1989). The geographic extent of mass coral spawning: clues to proximate and ultimate causal factors. Proc. 6th Int. Coral Reef Symp., Townsville (in press)

Pearse, J. S. (1968). Patterns of reproduction in four species of Indo-Pacific echinoderms. Proc. Indian Acad. Sci. 67: 247-267

Reed, S. A. (1971). Techniques for raising the planula larvae and newly settled polyps of *Pocillopora damicornis*. In: Lenhoff, H. M., Muscatine, L., Davis, L. V. (eds.) Experi-

mental coelenterate biology. Univ. Hawaii Press Richmond, R. H. (1981). Energetic considerations in the dispersal of *Pocillopora damicornis* planulae. Proc. 4th Int. Coral Reef Symp., Manila 2: 153–156

Richmond, R. H. (1985). Variations in the population biology of *Pocillopora damicornis* across the Pacific Ocean. Proc. 5th Int. Coral Reef Cong., Tahiti 6: 101–106

Richmond, R. H. (1987 a). Energetic relationships and biogeographical differences among fecundity, growth, and reproduction in the reef coral, *Pocillopora damicornis*. Bull. mar Sci. 41 (2): 595–604

Richmond, R. H. (1987b). Energetics, competency, and longdistance dispersal of planula larvae of the coral *Pocillopora* damicornis. Mar. Biol. 93: 527–533

Richmond, R. H. (1989). Competency and dispersal of spawned versus brooded coral planula larvae. Proc. 6th Int. Coral Reef Cong., Townsville (in press)

Richmond, R. H., Jokiel, P. L. (1984). Lunar periodicity in larva release in the reef coral *Pocillopora damicornis* at Enewetak and Hawaii. Bull. mar. Sci. 34: 280–287

Rinkevich, B., Loya, Y. (1979a). The reproduction of the Red Sea coral *Stylophora pistillata*. I. Gonads and planulae. Mar. Ecol. Prog. Ser. 1: 133–144

Rinkevich, B., Loya, Y. (1979b). The reproduction of the Red Sea coral Stylophora pistillata. II. Synchronization in breeding and seasonality of planula shedding. Mar. Ecol. Prog. Ser. 1: 145–152

- Rosen, B. R., Taylor, J. D. (1969). Reef coral from Aldabra: new mode of reproduction. Science 166: 119–120
- Rylaarsdam, K. W. (1983). Life histories and the abundance patterns of colonial corals on Jamaican reefs. Mar. Ecol. Prog. Ser. 13: 249–260
- Sammarco, P. W. (1982). Polyp bail-out: an escape response to environmental stress and a new means of reproduction in corals. Mar. Ecol. Prog. Ser. 10: 57–65
- Sammarco, P. W. (1985). The Great Barrier Reef vs. the Caribbean: comparisons of grazers, coral recruitment patterns and reef recovery. Proc. 5th Int. Coral Reef Cong., Tahiti 4: 391–397
- Sammarco, P. W. (1987). A comparison of ecological processes on coral reefs of the Caribbean and the Great Barrier Reef. In: Birkeland, C. (ed.) Comparison between Atlantic and Pacific tropical marine coastal ecosystems: community structure, ecological processes, and productivity. Univ. So. Pacif., Suva, Fiji, 1986. UNESCO Rep. mar. Sci. 46: 127–166
- Sammarco, P. W., Andrews, J. C. (1988). Localized dispersal and recruitment in Great Barrier Reef corals: The Helix Experiment. Science 239: 1422–1424
- Sammarco, P. W., Andrews, J. C. (1989). The Helix Experiment: differential localised dispersal and recruitment patterns in Great Barrier Reef corals. Limnol. Oceanogr. (in press)
- Sheppard, C. R. C. (1987). Coral species of the Indian Ocean and adjacent seas: a synomized compilation and some regional distribution patterns. Atoll Res. Bull. 307: 1–32
- Shlesinger, Y., Loya, Y. (1985). Coral community reproductive patterns: Red Sea versus the Great Barrier Reef. Science 228: 1333–1335
- Stimson, J. S. (1978). Mode and timing of reproduction in some common hermatypic corals of Hawaii and Enewetak. Mar. Biol. 48: 173–184
- Stoddart, J. A. (1983). The asexual production of planulae in the coral *Pocillopora damicornis*. Mar. Biol. 76: 279–284
- Stoddart, J. A. (1984). Genetical structure within populations of the coral *Pocillopora damicornis*. Mar. Biol. 81: 19–30
- Szmant, A. M. (1986). Reproductive ecology of Caribbean reef corals. Coral Reefs 5: 43–53
- Szmant-Froelich, A. (1984). Reef coral reproduction: diversity and community patterns. In: Advances in reef science.

This review was presented by Dr. P. W. Sammarco, Townsville, Australia

- Joint Meeting Intl. Soc. for Reef Studies and Atlantic Reef Committee. University of Miami, Miami, p. 122–123
- Szmant-Froelich, A., Reutter, M., Riggs, L. (1985). Sexual reproduction of *Favia fragum* (Esper): lunar patterns of gametogenesis, embryogenesis and planulation in Puerto Rico. Bull. mar. Sci. 37: 880–892
- Tomascik, T., Sander, F. (1987). Effects of eutrophication on reef-building corals. III. Reproduction of the reef-building coral *Porites porites*. Mar. Biol. 94: 77–94
- Tunnicliffe, V. (1981). Breakage and propagation of the stony coral Acropora cervicornis. Proc. natn. Acad. Sci. USA 78: 2427–2431
- Van Moorsel, G. W. N. M. (1983). Reproductive strategies in two closely related stony corals (*Agaricia*, Scleractinia). Mar. Ecol. Prog. Ser. 13: 273–283
- Vaughan, T. W. (1910). The recent Madreporaria of southern Florida. Yb. Carnegie Instn. Wash. 7: 135–144
- Veron, J. E. W. (1985). Corals of Australia and the Indo-Pacific. Angus and Robertson, London
- Veron, J. E. W., Wallace, C. C. (1984). Scleractinia of Eastern Australia, Part V, Family Acroporidae. A. I. M. S (Australian Institute of Marine Science, Townsville) Monogr. Ser. Vol. 6, p. 1–485
- Wallace, C. C. (1985 a). Four years of juvenile coral recruitment to five reef front sites. Proc. 5th Int. Coral Reef Cong., Tahiti 4: 385–390
- Wallace, C. C. (1985b). Reproduction, recruitment and fragmentation in nine sympatric species of the coral genus Acropora. Mar. Biol. 88: 217–233
- Williams, G. C. (1975). Sex and evolution. Princeton University Press, New Jersey
- Willis, B. L., Babcock, R. C., Harrison, P. L., Oliver, T. K. (1985). Patterns in the mass spawning of corals on the Great Barrier Reef from 1981 to 1984. Proc. 5th Int. Coral Reef Cong., Tahiti 4: 343–348
- Wyers, S. C. (1985). Sexual reproduction of the coral *Diploria strigosa* (Scleractinia, Favidae) in Bermuda: research in progress. Proc. 5th Int. Coral Reef Cong., Tahiti 4: 301–306
- Yamazato, K., Oshiro, M., Oshiro E. (1975). Reproductive biology of a scleractinian coral, Goniopora queenslandiae decima. Proc. 13th Pac. Sci. Congr. 1: 135 (Abstr.)

Manuscript first received: January 8, 1988 Revised version accepted: October 18, 1989